



US FedEx Shipping Instructions

- 1) Ship specimens Monday through Friday. If you complete the collection on the weekend, hold the specimen until the following Monday to ship.
- 2) Place the cardboard kit box containing the specimens and the completed test requisition form into the FedEx Clinical Pak.
- 3) Affix the provided Prepaid Return Label onto the bag in the designated location. Record the tracking number printed on the label if you want to track this delivery.
- 4) To schedule a pickup, call FedEx toll-free at 1-800-463-3339 (1-800-GO FEDEX). When you hear the automated greeting, say "Schedule a pick up." The next prompt asks if the shipment is more than 150lbs, say "No." At the next prompt, say "Schedule a Pick up with a Label or Stamp." Reply "YES" when asked if the word "Stamp" is written on the waybill. You will then be asked for your address information.

Do not use a drop box. FedEx will not accept specimens placed into a drop box.

If you are located in the United States and choose to use a different courier or level of service than provided or are located outside the United States, you must make your own shipping arrangements at your own expense.



US & CANADA: 1.800.323.2784 - UK: 0871.218.0052
ELSEWHERE: 1.630.377.8139 - FAX: 1.630.587.7860
3755 Illinois Avenue, St. Charles, IL USA 60174-2420
inquiries@doctorsdata.com - www.doctorsdata.com

GI360™ Essentials

Before You Start:

Please read all of the instructions carefully before beginning. Consult your physician for specific instructions and before stopping any medications. *Never discontinue prescription medications without first consulting your physician.*

Discontinue the Following:

Two Weeks Before The Test: Antibiotics, antiparasitics, antifungals, or probiotic supplements, proton pump inhibitors (PPIs) and Bismuth.

Two Days Before The Test: Aspirin and other NSAIDs, digestive enzymes, laxatives (particularly mineral oil, castor oil, and glycerin enemas/suppositories), activated charcoal, betaine HCl, antacids or bentonite clay.

Other Information:

- ◆ Do not collect samples when there is active bleeding from hemorrhoids or menstruation
- ◆ Wait at least 4 weeks from a colonoscopy or barium enema before collecting
- ◆ Do not contaminate the stool with urine or water

Verify Kit Contents:

- 1 Test requisition form
- 1 FedEx Clinical Pak with Prepaid Return Label
- 3 Collection containers (nested together; separate before use)

Day One Bag:	Day Two Bag:	Day Three Bag:
1 Black-capped vial	1 Black-capped vial	1 Black-capped vial
1 Disposable glove	1 Disposable glove	1 Yellow-capped vial
1 Absorbent material	1 Absorbent material	1 Disposable glove
		1 Absorbent material

- ◆ If you are missing kit components, please call Doctor's Data's Customer Service department for assistance.
- ◆ The Black-capped and Yellow-capped vials are filled to the middle of the vial with a liquid preservative. **If the liquid levels fall below the middle of the vial or there is evidence of leakage, CALL Customer Service for assistance.**
- ◆ Save the cardboard collection kit box to ship your samples back to the laboratory. Leave the absorbent material in the zip lock bag. Do not put it in the specimen vial.


Caution: Avoid contact of the skin and eyes with the fluid in the vials. For eye contact, flush with water thoroughly for 15 minutes. For skin contact, wash thoroughly with soap and water. For accidental ingestion, contact your physician immediately.

Collection One:

Use the contents of the Day One bag.




1) Collect your first stool specimen into the collection container. *DO NOT contaminate the specimen with either urine or water from the toilet.*



2) Unscrew the cap on the black-capped vial and, using the attached spoon, transport stool specimen into the vial. Take multiple portions from different areas of the collection container. Add stool until the total volume of stool and liquid reaches the fill line. **DO NOT OVERFILL.** Screw the cap on *tightly*.



3) Shake the black-capped vial vigorously for approximately 30 seconds to mix the stool specimen with the preservative in the vial.



4) Write the patient's name, the date of collection, and patient's date of birth on the specimen vial. (You do not have to provide an ID#. The lab will assign one upon arrival). ***The test cannot be performed without the patient information on the vial.***




5) Place the black-capped vial in the Day One zip-lock bag, then place the bag into the cardboard shipping box. Leave the box at room temperature.

Collection Two:

Use the contents of the Day Two bag.



6) Collect your second stool specimen into the collection container. *DO NOT contaminate the specimen with either urine or water from the toilet.*



7) Unscrew the cap on the black-capped vial and, using the attached spoon, transport stool specimen into the vial. Take multiple portions from different areas of the collection container. Add stool until the total volume of stool and liquid reaches the fill line. **DO NOT OVERFILL.** Screw the cap on *tightly*.



8) Shake the black-capped vial vigorously for approximately 30 seconds to mix the stool specimens with the preservative in the vial.



9) Write the patient's name, the date of collection, and patient's date of birth on the specimen vial. (You do not have to provide an ID#. The lab will assign one upon arrival). ***The test cannot be performed without the patient information on the vial.***



10) Place the black-capped vial into the Day Two zip-lock bag, then place the bag into the cardboard shipping box. Leave the box at room temperature.

Collection Three:


Use the contents of the Day Three bag.




11) Collect your third stool specimen into the collection container. Remember not to contaminate the specimen with urine or water from the toilet.




12) Unscrew the cap on the yellow-capped vial and, using the attached spoon, transport stool specimen into the vial. Take multiple portions from different areas of the collection container. Add stool until the total volume of stool and liquid reaches the fill line. **DO NOT OVERFILL.** Screw the cap on *tightly*.



13) Unscrew the cap on the black-capped vial and, using the attached spoon, transport stool specimen into the vial. Take multiple portions from different areas of the collection container. Add stool until the total volume of stool and liquid reaches the fill line. **DO NOT OVERFILL.** Screw the cap on *tightly*.



14) Shake both the yellow and black-capped vials vigorously for approximately 30 seconds to mix the stool specimens with the preservatives in the vials.



15) Write the patient's name, the date of collection, and patient's date of birth on both specimen vials. (You do not have to provide an ID#. The lab will assign one upon arrival). ***The test cannot be performed without the patient information on the vials.***



16) Place both the yellow and black-capped vials into the Day Three zip-lock bag, then place the bag into the cardboard shipping box. Check to be sure that all of the bags and vials are in the box.



17) Fill out the test requisition form completely and sign it. ***The test cannot be performed without the patient information on the vial and a properly filled out requisition.*** Place the form in the cardboard shipping box. The specimens are now ready for shipment.

To obtain the best results, Doctor's Data recommends shipping specimens as soon as possible. If you cannot ship the specimens the same day you finish the collection, such as over a weekend or holiday, place the cardboard box containing the yellow and black-capped vials into a refrigerator until you are ready to ship the specimens.

Consult your physician if you have any questions during the test collection process.